

# Relapsed ALL: Does End-Induction MRD Predict 1 yr or 2 yr EFS ?

Intensive Induction Therapy for Children with Acute  
Lymphoblastic Leukemia who Experience a Bone Marrow  
Relapse: Results from COG Study AALL01P2




**MRD as a Surrogate Endpoint Workshop**

**April 18, 2012**

Elizabeth A. Raetz, MD

William L. Carroll, MD

# Introduction of New Agents at First Relapse




# COG AALL01P2: Objectives and Eligibility

- **Primary Aim**
  - To develop a safe and active chemotherapy re-induction platform regimen for relapsed ALL, which could be used to evaluate novel agents in future trials
- **Eligibility**
  - Children 1-21 years of age with an initial ALL relapse involving the marrow (with or without associated extramedullary disease)

# COG AALL01P2: Patient Characteristics

- Timing of Relapse
  - 69 Early relapse (< 36 months from diagnosis)
  - 55 Late relapse ( $\geq$  36 months from diagnosis)
  - 4 Ph+
- Immunophenotype
  - 117 B lymphoblastic
  - 7 T-ALL
- Sites of Disease
  - 110 Isolated marrow, or marrow + other non-CNS EM site
  - 14 Marrow + CNS

# *AALL01P2 Study Design*



# COG AALL01P2: Accrual

## *Accrual*

- Study opened January 27, 2003
- Permanently closed December 30, 2005
- 124 eligible patients enrolled after first amendment/therapy modification

# COG AALL01P2: Remission Re-induction Rates

	<b><i>Very Early Relapse</i></b> ( $< 18$ mo from dx)	<b><i>Early Relapse</i></b> ( $< 36$ mo from dx)	<b><i>Late Relapse</i></b> ( $\geq 36$ mo from dx)
All patients	45.0% $\pm$ 11.1% (n=24)	68.3% $\pm$ 5.9% (n=69)	96.3% $\pm$ 2.6% (n=55)
B-lineage, CNS-, Ph-	46.2% $\pm$ 13.8% (n=16)	72.0% $\pm$ 6.4% (n=55)	95.5% $\pm$ 3.1% (n=45)

# MRD Response Rates According to Timing of Relapse

	MRD Positive Patients * (%)	
Treatment Block	Early Relapse	Late Relapse
1	75 ± 7 (n = 36)	51 ± 8 (n = 43)
2	70 ± 10 (n = 23)	41 ± 4 (n = 32)
3	43 ± 11 (n = 21)	25 ± 8 (n = 28)


\* All patients in complete remission at the end of block 1

At the end of three blocks of therapy, 16 (33%) of 49 of patients overall, remained MRD positive




# COG AALL01P2: EFS and OS

All B Lymphoblastic Patients




All B Lymphoblastic Patients



*Data updated 2/24/12, Xiaomin Lu PhD*

# COG AALL01P2: Outcomes According to Time to Relapse


## B Lymphoblastic Patients




Data updated 2/24/12, Xiaomin Lu PhD

# Outcomes According to Block 1 MRD Response


## All B lymphoblastic patients with CR by MRD (0.01%)




## Early relapse B lymphoblastic with CR by MRD (0.01%)



## Late relapse B lymphoblastic with CR by MRD (0.01%)



# Kinetics of MRD Response



~ 70% of patients who achieve CR2, but have detectable MRD at the end of block 1, will have sustained remissions and further reductions in MRD with ongoing therapy

# **Reinduction Chemoimmunotherapy with Epratuzumab in Relapsed ALL in Children, Adolescents and Young Adults: COG Study ADVL04P2**

# COG ADVL04P2: Objectives and Eligibility

- Primary Aim
  - To determine if addition of epratuzumab, an anti-CD22 monoclonal antibody, to the AALL01P2 backbone chemotherapy regimen improves rates of second complete remission (CR2) in children and young adults with ALL and early bone marrow relapse (<36 months from initial diagnosis)
- Eligibility
  - Children and young adults ages 2-30 years with first, early marrow relapse of CD22+ B-cell precursor ALL, with or without extramedullary disease

# Study Design

## Part A

Feasibility

Reduction Phase:

Day -14 to 0



Block 1



Triple Induction\*

Block 2

Block 3

Response

CR2 rate &  
MRD

Triple Induction\*

Block 1



Block 2

Block 3

CR2 rate &  
MRD

↑ = Epratuzumab dose (360 mg/m<sup>2</sup>)

## Part B

Phase 2 Pilot:

*B1 Cohort*

*B2 Cohort*

# Accrual

- Part A (Feasibility): Epratuzumab twice weekly x 4, then weekly x 4
  - 15 patients from 1/06-6/06
- Part B1 (Phase 2): Epratuzumab weekly x 4
  - 56 patients (54 eligible) from 1/07-12/08
- Part B2 (Phase 2): Epratuzumab twice weekly x 8
  - 60 patients from 1/09-1/11



# Results Part A\*: Reduction Phase and Block 1 Responses

Response	Number of Patients (n=15)
<i>Reduction Phase</i>	
• Stable Disease	8
• Minimal Remission Cytolytic	4
• Progressive Disease	3
<i>End of Block 1</i>	
• Complete Remission	9*
• Partial Remission (M2 marrow)	1
• Stable Disease	1
• Progressive Disease	1
• Died from infection or removed from protocol	3

**\* Seven of nine patients achieving second CR had no detectable MRD**

# Responses for B1 and B2 Cohorts

	B1 Cohort (weekly x 4)	B2 Cohort (twice weekly x 8)
Eligible Patients	54	60
Very Early Relapse (< 18mo)	23	19
Median Age at Relapse (yrs)	10.2	8.4
Extramedullary Disease	3	9
Response Evaluable Patients	48	50
End Block 1 CR2 Rate	65% (31/48)	66% (33/50)
End Block 1 MRD Available	31	31
End Block 1 MRD Neg (<0.01%)	45% (14/31)	39% (12/31)
End Block 1 MRD Neg Pooled	42% (26/62)*	

**\*Significantly higher than the 25% (9/36) with chemotherapy alone on AALL01P2 (one-sided p=0.001)**

# Conclusions

- Rates of MRD positivity are much higher in relapsed ALL than newly diagnosed disease
- Early MRD response was a strong predictor for EFS on the COG AALL01P2 trial
- The kinetic pattern of MRD was also predictive of longer term outcomes in relapsed ALL on COG AALL01P2
- The combination of MRD response and timing of relapse identified different risk groups of patients, suggesting that MRD may be helpful in stratifying salvage therapy

# Acknowledgements

Peter Adamson

Susan Blaney

Michael Borowitz

Mitch Cairo

Bill Carroll

Mini Devidas

Paul Gaynon

David Goldenberg

Steve Hunger

Mignon Loh

Xiaomin Lu

Gregory Reaman

Jim Whitlock

Naomi Winick

Charlotte Wood